Scientific name: Berberis vulgaris USDA Plants Code: BEVU Common names: Common barberry Southern Europe and Asia Native distribution: March 4, 2008; edited 7 April 2009 Date assessed: Jinshuang Ma, Gerry Moore Assessors: Reviewers: LIISMA SRC 9-24-2008 Form version date: 22 October 2008 Date Approved:

**New York Invasiveness Rank:** Moderate (Relative Maximum Score 50.00-69.99)

Distribution and Invasiveness Rank (Obtain from PRISM invasiveness ranking form)			
			PRISM
	Status of this species in each PRISM:	Current Distribution	Invasiveness Rank
1	Adirondack Park Invasive Program	Not Assessed	Not Assessed
2	Capital/Mohawk	Not Assessed	Not Assessed
3	Catskill Regional Invasive Species Partnership	Not Assessed	Not Assessed
4	Finger Lakes	Not Assessed	Not Assessed
5	Long Island Invasive Species Management Area	Common	Moderate
6	Lower Hudson	Not Assessed	Not Assessed
7	Saint Lawrence/Eastern Lake Ontario	Not Assessed	Not Assessed
8	Western New York	Not Assessed	Not Assessed

Inv	asiveness Ranking Summary	Total (Total Answered*)	Total
(see	details under appropriate sub-section)	Possible	
1	Ecological impact	40 (20)	6
2	Biological characteristic and dispersal ability	25 ( <u>25</u> )	23
3	Ecological amplitude and distribution	25 ( <u>25</u> )	21
4	Difficulty of control	10 ( <u>10</u> )	5
	Outcome score	100 ( <u>80</u> ) <sup>b</sup>	55 <sup>a</sup>
	Relative maximum score †		68.75
	New York Invasiveness Rank §	Moderate (Relative Maximum Score 50.00-69.99)	

<sup>\*</sup> For questions answered "unknown" do not include point value in "Total Answered Points Possible." If "Total Answered Points Possible" is less than 70.00 points, then the overall invasive rank should be listed as "Unknown." †Calculated as 100(a/b) to two decimal places.

\$Very High >80.00; High 70.00-80.00; Moderate 50.00-69.99; Low 40.00-49.99; Insignificant <40.00

#### A. DISTRIBUTION (KNOWN/POTENTIAL): Summarized from individual PRISM forms

	s this species been documented to persist without in NY? (reliable source; voucher not required)	Partnerships for Regional Invasive Species Management
$\boxtimes$	Yes – continue to A1.2	2008
	No – continue to A2.1	APIPP
A1.2. In v	which PRISMs is it known (see inset map)?	SLELO
$\boxtimes$	Adirondack Park Invasive Program	Capital
$\boxtimes$	Capital/Mohawk	Finger Lakes Mohawk
$\boxtimes$	Catskill Regional Invasive Species Partnership	Western NY Ones
	Finger Lakes	CRISP
$\boxtimes$	Long Island Invasive Species Management Area	Lower
$\boxtimes$	Lower Hudson	Hudson
	Saint Lawrence/Eastern Lake Ontario	Liisma
	Western New York	San

	Documentation: Sources of information: Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008 A2.1. What is the likelihood that this species will occur		of cultivation given the climate in		
Not A	the following PRISMs? (obtain from PRISM invasive no Assessed Adirondack Park Invasive Program		reactivation given the elimate in		
Not A	Assessed Capital/Mohawk				
	Assessed Catskill Regional Invasive Species P	artnership			
	Assessed Finger Lakes				
-	Likely Long Island Invasive Species Manag Assessed Lower Hudson	ement Area			
	Assessed Lower Hudson Assessed Saint Lawrence/Eastern Lake Ontario	2			
	Assessed Western New York	)			
1011	Documentation:				
	Sources of information (e.g.: distribution models, literat Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008		:		
f th	e species does not occur and is not likely to oc	cur with any of t	he PRISMs, then stop here		
	as there is no need to as	sess the species.			
	A2.2. What is the current distribution of the species in e	ach PRISM? (obtain	rank from PRISM invasiveness		
	ranking forms)		Distribution		
	Adirondack Park Invasive Program		Not Assessed		
	Capital/Mohawk		Not Assessed		
	Catskill Regional Invasive Species Partnership		Not Assessed		
	Finger Lakes		Not Assessed		
	Long Island Invasive Species Management Area		Common		
	Lower Hudson		Not Assessed		
	Saint Lawrence/Eastern Lake Ontario		Not Assessed		
	Western New York		Not Assessed		
	Documentation:				
	Sources of information:				
	Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008	•			
	A2.3. Describe the potential or known suitable habitats habitats not under active human management.				
	Aquatic Habitats Wetland Habitats	Uı	pland Habitats		
	Salt/brackish waters Salt/brackish		Cultivated*		
	☐ Freshwater tidal ☐ Freshwater ☐ Rivers/streams ☐ Peatlands	marshes L	<ul><li>✓ Grasslands/old fields</li><li>✓ Shrublands</li></ul>		
	☐ Natural lakes and ponds ☐ Shrub swan	nns	∑ Situations  X Forests/woodlands		
		etlands/riparian [	Alpine		
	☐ Reservoirs/impoundments* ☐ Ditches*		Roadsides*		
	Other potential or known suitable habitats within New Y	/or coastal dunes ork:			
	Documentation:				
	Sources of information:				
	Kern 1921a: Mayhury 2005: Brooklyn Rotanic Garden 2008: author's (Moore's) personal observations				

### **New York** NON-NATIVE PLANT INVASIVENESS RANKING FORM

### **B. INVASIVENESS RANKING**

1	<b>ECOLOGICAL</b>	<b>IMPACT</b>
	LCOLOGICAL	IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII

regime,	pact on Natural Ecosystem Processes and System-Wide Parameters (e.g. fire geomorphological changes (erosion, sedimentation rates), hydrologic regime, and mineral dynamics, light availability, salinity, pH)  No perceivable impact on ecosystem processes based on research studies, or the absence of impact information if a species is widespread (>10 occurrences in minimally managed areas), has been well-studied (>10 reports/publications), and has been present in the	0
B.	northeast for >100 years. Influences ecosystem processes to a minor degree (e.g., has a perceivable but mild influence on soil nutrient availability)	3
C.	Significant alteration of ecosystem processes (e.g., increases sedimentation rates along streams or coastlines, reduces open water that are important to waterfowl)	7
D.	Major, possibly irreversible, alteration or disruption of ecosystem processes (e.g., the species alters geomorphology and/or hydrology, affects fire frequency, alters soil pH, or fixes substantial levels of nitrogen in the soil making soil unlikely to support certain native plants or more likely to favor non-native species)	10
U.	Unknown Score	U
1.2. Imp A. B. C.	Documentation:  Identify ecosystem processes impacted (or if applicable, justify choosing answer A in the absence of impact information)  While the species has been present since the 1800s, ecological studies on its impact to ecosystem processes and parameters is largely lacking, with most of the early research when the plant was probably more prevalent focusing on the species' distribution and its effects on cereal grains (it serves as the alternate host for a rust fungus that infects grasses). Answer A was not given because of the lack of ecological studies within the B. vulgaris literature. Question was not answered because of lack of ecological studies. Sources of information:  Kern, 1921a; Maybury, 2005.  Coact on Natural Community Structure  No perceived impact; establishes in an existing layer without influencing its structure Influences structure in one layer (e.g., changes the density of one layer)  Significant impact in at least one layer (e.g., creation of a new layer or elimination of an	0 3 7
	existing layer)  Major alteration of structure (e.g., covers canopy, eradicating most or all layers below)	·
D. U.	Unknown	10
	Score	3
	Documentation: Identify type of impact or alteration: Currently the species is only occasionally noted in the shrub layer where it can increase the density in that layer. Kern (1921a) reported "thousands of bushes, forming thickets in pastures" (and thus creating a new layer) from Pennsylvania in the early 1900s but recent evidence (Maybury, 2005; Moore's personal observations) do not show this. Sources of information: Kern, 1921a; Maybury, 2005; author's (Moore's) personal observations.	
	pact on Natural Community Composition  No perceived impact; causes no apparent change in native populations	0
A. B.	Influences community composition (e.g., reduces the number of individuals in one or more	0 3
C.	native species in the community) Significantly alters community composition (e.g., produces a significant reduction in the	7

D. U.	population size of one or more native species in the community) Causes major alteration in community composition (e.g., results in the extirpation of one or several native species, reducing biodiversity or change the community composition towards species exotic to the natural community) Unknown	10
	Score	3
	Documentation: Identify type of impact or alteration: Has recently been noted to form small thickets where it reduced the number of native species present. Sources of information: Maybury, 2005; author's (Moore's) personal observations.	
the anin Exampl connect soil/sed native s	pact on other species or species groups (cumulative impact of this species on nals, fungi, microbes, and other organisms in the community it invades. es include reduction in nesting/foraging sites; reduction in habitat ivity; injurious components such as spines, thorns, burrs, toxins; suppresses iment microflora; interferes with native pollinators and/or pollination of a pecies; hybridizes with a native species; hosts a non-native disease which a native species)	
A.	Negligible perceived impact	0
В.	Minor impact	3
C.	Moderate impact	7
D.	Severe impact on other species or species groups	10
U.	Unknown	
	Score	U
	Documentation:  Identify type of impact or alteration:  Serves as the alternate host for the rust Puccinia graminis, which, besides cultivated cereal grasses, can infect many species of native grass genera (e.g., Ammophila, Buchloe, Calamagrostis, Danthonia Deschampsia, Distichilis, Glyceria, Muhlenbergia, Panicum, Poa, Puccinellia, Sphenopholis, Sporobolus, Stipa) (Farr et al, 1989; University of Nebraska, 2003; Leonard & Szabo, 2005). Berberis thunbergii has not been reported as a host for P. graminis, and the native Berberis canadensis, which can serve as the host, does not occur in New York (Weldy & Werier, 2005). Therefore, B. vulgaris is the only known alternate host in New York for P. graminis. However, B. vulgaris is only occasionaly reported from the area (Maybury, 2005; Moore's personal obervations) and observations of the rust fungus on Berberis and native grass species are lacking (Moore personal observations). Has thorns but not sure of impact on other species.  Sources of information:  Farr et al., 1989; University of Nebraska, 2003; Leonard & Szabo, 2005, Maybury, 2005; Weldy & Werier, 2005; author's (Moore's) personal obervations.  Total Possible	20
	Section One Total	6
	OLOGICAL CHARACTERISTICS AND DISPERSAL ABILITY	
2.1. Mo A.	de and rate of reproduction (provisional thresholds, more investigation needed)  No reproduction by seeds or vegetative propagules (i.e. plant sterile with no sexual or assayual reproduction)	0
В.	asexual reproduction).  Limited reproduction (fewer than 10 viable seeds per plant AND no vegetative reproduction; if viability is not known, then maximum seed production is less than 100 seeds per plant and no vegetative reproduction)	1

C.	Moderate reproduction (fewer than 100 viable seeds per plant - if viability is not known, then maximum seed production is less than 1000 seeds per plant - OR limited successful vegetative spread documented)	2
D.	Abundant reproduction with vegetative asexual spread documented as one of the plants prime reproductive means OR more than 100 viable seeds per plant (if viability is not known, then maximum seed production reported to be greater than 1000 seeds per plant.)	4
U.	Unknown	
	Score	4
	Documentation:	
	Describe key reproductive characteristics (including seeds per plant): Individuals can produce fruits (which are 1-few seeded) in the low hundreds and viability of the seeds in the species (Morinaga, 1926) and genus in general (Kern, 1921; Davis, 1927; Allen & Wilson, 1992) can be (above 70%). Sources of information:	
	Kern, 1921a; Morinaga, T. 1926; Davis, 1927; Allen & Wilson, 1992.	
	ate potential for long-distance dispersal (e.g. bird dispersal, sticks to animal hair,	
•	fruits, pappus for wind-dispersal)  Does not occur (no long-distance dispersal mechanisms)	0
A. B.	Infrequent or inefficient long-distance dispersal (occurs occasionally despite lack of	0 1
Ъ.	adaptations)	1
C.	Moderate opportunities for long-distance dispersal (adaptations exist for long-distance dispersal, but studies report that 95% of seeds land within 100 meters of the parent plant)	2
D.	Numerous opportunities for long-distance dispersal (adaptations exist for long-distance dispersal and evidence that many seeds disperse greater than 100 meters from the parent	4
	plant)	
U.	Unknown	
	Score	4
	Documentation: Identify dispersal mechanisms:	
	Fruits are bird and mammal dispersed, including cattle (at least in earlier times; Kern 1921b).	
	Sources of information:	
	Kern, 1921a, 1921b; Mehrhoff et al., 2003; Maybury, 2005; author's (Moore's) personal observations.	
	tential to be spread by human activities (both directly and indirectly – possible	
	nisms include: commercial sales, use as forage/revegetation, spread along	
•	ys, transport on boats, contaminated compost, land and vegetation	
manage A.	ement equipment such as mowers and excavators, etc.)  Does not occur	0
В.	Low (human dispersal to new areas occurs almost exclusively by direct means and is	1
	infrequent or inefficient)	•
C.	Moderate (human dispersal to new areas occurs by direct and indirect means to a moderate extent)	2
D.	High (opportunities for human dispersal to new areas by direct and indirect means are numerous, frequent, and successful)	3
U.	Unknown	
	Score	1
	Documentation:	
	Identify dispersal mechanisms: Plant is not sold and dispersal by direct means does not seem likely.	
	Sources of information:	
2.4.01	Maybury, 2005; J. Lehrer, pers. comm	
2.4. Cn	aracteristics that increase competitive advantage, such as shade tolerance,	

5

ability to grow on infertile soils, perennial habit, fast growth, nitrogen fixation, allelopathy, etc. A. Possesses no characteristics that increase competitive advantage 0 Possesses one characteristic that increases competitive advantage B. 3 Possesses two or more characteristics that increase competitive advantage C. 6 U. Unknown Score 6 Documentation: Evidence of competitive ability: Shade tolerant, perennial, can grow on infertile soils. Sources of information: Kern, 1921a, 1921b; Maybury, 2005; J. Lehrer pers. comm.; C. Scheer pers. comm.. 2.5. Growth vigor Does not form thickets or have a climbing or smothering growth habit 0 A. Has climbing or smothering growth habit, forms a dense layer above shorter vegetation, 2 forms dense thickets, or forms a dense floating mat in aquatic systems where it smothers other vegetation or organisms Unknown U. Score 2 Documentation: Describe growth form: Observed to form small thickets; earlier literature reported it forming large thickets. Sources of information: Kern, 1921; author's (Moore's) personal observations. 2.6. Germination/Regeneration Requires open soil or water and disturbance for seed germination, or regeneration from 0 vegetative propagules. Can germinate/regenerate in vegetated areas but in a narrow range or in special conditions B. 2 Can germinate/regenerate in existing vegetation in a wide range of conditions C. 3 U. Unknown (No studies have been completed) 3 Score Documentation: Describe germination requirements: Observed to germinate in exisiting vegetative conditions. Sources of information: Kern 1921a, 1921b, Maybury, 2005; author's (Moore's) personal observations. 2.7. Other species in the genus invasive in New York or elsewhere No Α. 0 Yes 3 B. U. Unknown Score 3 Documentation: Species: Berberis thunbergii. **Total Possible** Section Two Total 23

#### 3. ECOLOGICAL AMPLITUDE AND DISTRIBUTION

3.1. Density of stands in natural areas in the northeastern USA and eastern Canada

(use same definition as Gleason & Cronquist which is: "The part of the United States covered extends from the Atlantic Ocean west to the western boundaries of Minnesota, Iowa, northern Missouri, and southern Illinois, south to the southern boundaries of Virginia, Kentucky, and Illinois, and south to the Missouri River in Missouri. In Canada the area covered includes Nova Scotia, Prince Edward Island, New Brunswick, and parts of Quebec and Ontario lying south of the 47th parallel of latitude")

iatitude		
A.	No large stands (no areas greater than 1/4 acre or 1000 square meters)	0
B.	Large dense stands present in areas with numerous invasive species already present or disturbed landscapes	2
C.	Large dense stands present in areas with few other invasive species present (i.e. ability to invade relatively pristine natural areas)	4
U.	Unknown	
	Score	2
	Documentation:	
	Identify reason for selection, or evidence of weedy history: Some large populations are reported in NYS, including in Bergen Byron Swamp, Genesee Co., western NY. It is not a dense stand, but covers >1/4 acre. Most populations in NYS are small, just a few plants (D. Werier). Some, but not all, populations are in disturbed areas, but more information is needed on size and disturbance. Large populations were reported in earlier times from Pennsylvania and elsewhere (e.g., Kern, 1921a; Mack, 2003).	
	Sources of information: Kern, 1921a; Mack, 2003; Maybury, 2005; author's (Moore's) and D. Werier's personal observations.	
3.2. Nu	mber of habitats the species may invade	
A.	Not known to invade any natural habitats given at A2.3	0
В.	Known to occur in two or more of the habitats given at A2.3, with at least one a natural habitat.	1
C.	Known to occur in three or more of the habitats given at A2.3, with at least two a natural habitat.	2
D.	Known to occur in four or more of the habitats given at A2.3, with at least three a natural habitat.	4
E.	Known to occur in more than four of the habitats given at A2.3, with at least four a natural habitat.	6
U.	Unknown	
	Score	6
	Documentation:	
	Identify type of habitats where it occurs and degree/type of impacts: See A2.3.	
	Sources of information: Kern, 1921a; Maybury, 2005; Brooklyn Botanic Garden 2008; author's (Moore's) personal observations.	
3.3. Ro	le of disturbance in establishment	
A.	Requires anthropogenic disturbances to establish.	0
B.	May occasionally establish in undisturbed areas but can readily establish in areas with natural or anthropogenic disturbances.	2
C.	Can establish independent of any known natural or anthropogenic disturbances.	4
U.	Unknown	
	Score	2
	Documentation:	
	Identify type of disturbance:	

	Usually found in disturbed habitats, it has also been noted in undisturbed areas.  Sources of information:  Mehrhoff et al., 2003; Maybury, 2005; author's (Moore's) and D. Werier's personal observations.		
3.4. Cli	mate in native range		
A.	Native range does not include climates similar to New York		0
В.	Native range possibly includes climates similar to at least part of New York.		1
C.	Native range includes climates similar to those in New York		3
U.	Unknown		3
0.	Scor	re	3
	Documentation:		<u>J</u>
	Describe what part of the native range is similar in climate to New York:		
	Europe and temperate Asia.		
	Sources of information:		
	Whittemore, 1997; Brooklyn Botanic Garden, 2008.		
3.5. Cu	rrent introduced distribution in the northeastern USA and eastern Canada (see	9	
questio	n 3.1 for definition of geographic scope )		
A.	Not known from the northeastern US and adjacent Canada		0
B.	Present as a non-native in one northeastern USA state and/or eastern Canadian province.		1
C.	Present as a non-native in 2 or 3 northeastern USA states and/or eastern Canadian provinces.		2
D.	Present as a non-native in 4–8 northeastern USA states and/or eastern Canadian provinces,		3
D.	and/or categorized as a problem weed (e.g., "Noxious" or "Invasive") in 1 northeastern state		5
	or eastern Canadian province.		
E.	Present as a non-native in >8 northeastern USA states and/or eastern Canadian provinces.		4
	and/or categorized as a problem weed (e.g., "Noxious" or "Invasive") in 2 northeastern		
	states or eastern Canadian provinces.		
U.	Unknown		4
	Scor	re	4
	Documentation:		
	Identify states and provinces invaded:		
	CI DETATI IN MA MID ME MI MIN NH NI NY DH PA RI VA VI WI WV		
	CT, DE, IA, IL, IN, MA, MD, ME, MI, MN, NH, NJ, NY, OH, PA, RI, VA, VT, WI, WV;	;	
	NB, NF, NS, ON, PE, QC.	;	
	NB, NF, NS, ON, PE, QC. Sources of information:		
	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from		
	NB, NF, NS, ON, PE, QC. Sources of information:		
	<ul> <li>NB, NF, NS, ON, PE, QC.</li> <li>Sources of information:</li> <li>See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.</li> </ul>		
3.6. Cu	<ul> <li>NB, NF, NS, ON, PE, QC.</li> <li>Sources of information:</li> <li>See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.</li> </ul>		
	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  rrent introduced distribution of the species in natural areas in the eight New		
	<ul> <li>NB, NF, NS, ON, PE, QC.</li> <li>Sources of information: <ul> <li>See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.</li> </ul> </li> <li>U.S.D.A., 2008.</li> </ul>		0
York S	<ul> <li>NB, NF, NS, ON, PE, QC.</li> <li>Sources of information:         <ul> <li>See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.</li> </ul> </li> <li>U.S.D.A., 2008.</li> <li>rrent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)</li> </ul>		
York S A.	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  rrent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)  Present in none of the PRISMs		1
York S A. B. C.	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  rrent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)  Present in none of the PRISMs  Present in 1 PRISM		1 2
York S A. B. C. D.	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  rrent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)  Present in none of the PRISMs  Present in 1 PRISM  Present in 2 PRISMs		1 2 3
York S A. B. C. D. E.	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  Trent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)  Present in none of the PRISMs  Present in 1 PRISM  Present in 2 PRISMs  Present in 3 PRISMs		1 2
York S A. B. C. D.	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  rrent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)  Present in none of the PRISMs  Present in 1 PRISM  Present in 2 PRISMs  Present in 3 PRISMs  Present in more than 3 PRISMs or on the Federal noxious weed lists  Unknown	n	1 2 3 4
York S A. B. C. D. E.	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  rrent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)  Present in none of the PRISMs  Present in 1 PRISM  Present in 2 PRISMs  Present in 3 PRISMs  Present in more than 3 PRISMs or on the Federal noxious weed lists	n	1 2 3
York S A. B. C. D. E.	NB, NF, NS, ON, PE, QC. Sources of information:  • See known introduced range in plants.usda.gov, and update with information from states and Canadian provinces.  U.S.D.A., 2008.  rrent introduced distribution of the species in natural areas in the eight New tate PRISMs (Partnerships for Regional Invasive Species Management)  Present in none of the PRISMs  Present in 1 PRISM  Present in 2 PRISMs  Present in 3 PRISMs  Present in more than 3 PRISMs or on the Federal noxious weed lists  Unknown	n	1 2 3 4

All PRISMS; see A1.1. Sources of information: Weldy & Werier, 2005; Brooklyn Botanic Garden, 2008. **Total Possible** Section Three Total 21 4. DIFFICULTY OF CONTROL 4.1. Seed banks Seeds (or vegetative propagules) remain viable in soil for less than 1 year, or does not make A. 0 viable seeds or persistent propagules. Seeds (or vegetative propagules) remain viable in soil for at least 1 to 10 years 2 B. C. Seeds (or vegetative propagules) remain viable in soil for more than 10 years 3 U. Unknown Score 2 Documentation: Identify longevity of seed bank: Seeds can remain viable for over a year, but no evidence for 10 years. Sources of information: Morinaga, 1926; Allen & Wilson, 1992. 4.2. Vegetative regeneration No regrowth following removal of aboveground growth A. 0 Regrowth from ground-level meristems B. 1 Regrowth from extensive underground system C. 2 Any plant part is a viable propagule 3 D. U. Unknown Score 1 Documentation: Describe vegetative response: Regrowth from ground-level meristems. Sources of information: Maybury, 2005. 4.3. Level of effort required Management is not required: e.g., species does not persist without repeated anthropogenic 0 disturbance. Management is relatively easy and inexpensive: e.g. 10 or fewer person-hours of manual 2 effort (pulling, cutting and/or digging) can eradicate a 1 acre infestation in 1 year (infestation averages 50% cover or 1 plant/100 ft<sup>2</sup>). Management requires a major short-term investment: e.g. 100 or fewer person-hours/year of 3 manual effort, or up to 10 person-hours/year using mechanical equipment (chain saws, mowers, etc.) for 2-5 years to suppress a 1 acre infestation. Eradication is difficult, but possible (infestation as above). Management requires a major investment: e.g. more than 100 person-hours/year of manual 4 effort, or more than 10 person hours/year using mechanical equipment, or the use of herbicide, grazing animals, fire, etc. for more than 5 years to suppress a 1 acre infestation. Eradication may be impossible (infestation as above). U. Unknown Score 2

#### Documentation:

Identify types of control methods and time-term required:

Control can be effected through herbicide (e.g., glyphosphate), hand-pulling, and digging.

Known localities are mostly small, so management not expected to be difficult.

Sources of information:

Mehrhoff, 2003; Maybury, 2005; author's (Moore's) personal observations

Total Possible
Section Four Total

5

Total for 4 sections Possible
Total for 4 sections

55

#### C. STATUS OF CULTIVARS AND HYBRIDS:

At the present time (May 2008) there is no protocol or criteria for assessing the invasiveness of cultivars independent of the species to which they belong. Such a protocol is needed, and individuals with the appropriate expertise should address this issue in the future. Such a protocol will likely require data on cultivar fertility and identification in both experimental and natural settings.

Hybrids (crosses between different parent species) should be assessed individually and separately from the parent species wherever taxonomically possible, since their invasiveness may differ from that of the parent species. An exception should be made if the taxonomy of the species and hybrids are uncertain, and species and hybrids can not be clearly distinguished in the field. In such cases it is not feasible to distinguish species and hybrids, and they can only be assessed as a single unit.

Some cultivars of the species known to be available: purpurea, dulcis, possibly superba

#### **References for species assessment:**

Allen, R.B. and J.B. Wilson. 1992. Fruit and seed production in Berberis darwinii Hook., a shrub recently naturalised in New Zealand. New Zealand Journal of Botany 30: 45-55.

Brooklyn Botanic Garden. 2008. AILANTHUS database. [Accessed on March 4 2008.] Gleason, H. A. & A. Cronquist. 1991. Manual of Vascular Plants of Northeastern United States and Adjacent Canada. 2nd ed. The New York Botanical Garden, Bronx. 993 pp.

Davis, O. H. 1927. Germination and early growth of Cornus florida, Sambucus canadensis, and berberis thunbergii. Botanical Gazette 84(3): 225-263.

Farr, D.F., G.F. Bills, G.P. Chamuris, and A.Y. Rossman. Fungi of Plants and Plant Products in the United States. American Phytopathological Society. Saint Paul, Minnesota. 1252 pp.

Invasive Species Specialist Group 2005. Global Invasive Species Database < http://www.issg.org/database/species/ecology.asp?si=592&fr=1&sts=sss>. [Accessed March 4, 2008].

Kern, F.D. 1921a. Distribution of Berberis vulgaris in Pennsylvania. Bulletin of Torrey Botanical Club 48(10): 263-269.

Kern, F.D. 1921. Observations of the dissemination of the barberry. Ecology 2(3): 211-214.

Leonard, K.J. and L.J. Szabo. 2005. Stem rust of small grains and grasses caused by Puccinia graminis. Molecualr Plant Pathology 6(2): 99-111.

Mack, R.N. 2003. Plant naturalizations and invasions in the eastern United States. Annals of the Missouri Botanical Garden 90(1): 77-90.

Maybury, K. 2005. Berberis vulgaris. U.S. Invasive Species Impact Rank (I-Rank). NatureServe Explorer. <a href="https://www.natureserve.org">www.natureserve.org</a>. [Accessed on March 4, 2008.]

Mehrhoff, L.J., J.A. Silander, Jr., S.A. Leicht and E. Mosher. 2003. IPANE: Invasive Plant Atlas of New England. Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT. <a href="http://invasives.eeb.uconn.edu/ipane/">http://invasives.eeb.uconn.edu/ipane/</a>. [Accessed September 5, 2008.]

Morinaga, T. 1926. Effect of alternating temperatures upon the germination of seeds. American Journal of botany 60: 557-560.

United States Department of Agriculture. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, LA <a href="http://plants.usda.gov">http://plants.usda.gov</a>. [Accessed 4 March 2008.]

United States Department of Agriculture, National Resources Conservation Service. 2008. The PLANTS Database. National Plant Data Center, Baton Rouge, Louisiana. <plants.usda.gov>. [Accessed on March 4, 2008.]

University of Nebraska. 2003. Department of Plant Pathology Disease Descriptions. <nudistance.unl.edu/homer/disease/agron>. [Accessed March 4, 2008.]

Weldy, T. and D. Werier. 2005. New York Flora Atlas. [S.M. Landry, K.N. Campbell, and L.D. Mabe (original application development), Florida Center for Community Design and Research. University of South Florida]. New York Flora Association, Albany, New York. <atlas.nyflora.org/>. [Accessed on March 4, 2008.]

Whittemore, A. T. 1997. 3: 278. Berberis. In: Flora of North America Editorial Committee, eds. 1993+. Flora of North America North of Mexico. 12+ vols. New York and Oxford. Vol. 3, pp. 278-279.

**Citation:** This NY ranking form may be cited as: Jordan, M.J., G. Moore and T.W. Weldy. 2008. Invasiveness ranking system for non-native plants of New York. Unpublished. The Nature Conservancy, Cold Spring Harbor, NY; Brooklyn Botanic Garden, Brooklyn, NY; The Nature Conservancy, Albany, NY. Note that the order of authorship is alphabetical; all three authors contributed substantially to the development of this protocol.

Acknowledgments: The NY form incorporates components and approaches used in several other systems, cited in the references below. Valuable contributions by members of the Long Island Invasive Species Management Area's Scientific Review Committee were incorporated in revisions of this form. Original members of the LIISMA SRC included representatives of the Brooklyn Botanic Garden; The Nature Conservancy; New York Natural Heritage Program, New York Sea Grant; New York State Office of Parks, Recreation and Historic Preservation; National Park Service; Brookhaven National Laboratory; New York State Department of Environmental Conservation Region 1; Cornell Cooperative Extension of Suffolk/Nassau Counties; Long Island Nursery and Landscape Association; Long Island Farm Bureau; SUNY Farmingdale Ornamental Horticulture Department; Queens College Biology Department; Long Island Botanical Society; Long Island Weed Information Management System database manager; Suffolk County Department of Parks, Recreation and Conservation; Nassau County Department of Parks, Recreation and Museums; Suffolk County Soil & Water Conservation District.

### References for ranking form:

- Carlson, Matthew L., Irina V. Lapina, Michael Shephard, Jeffery S. Conn, Roseann Densmore, Page Spencer, Jeff Heys, Julie Riley, Jamie Nielsen. 2008. Invasiveness ranking system for non-native plants of Alaska. Technical Paper R10-TPXX, USDA Forest Service, Alaska Region, Anchorage, AK XX9. Alaska Weed Ranking Project may be viewed at: <a href="http://akweeds.uaa.alaska.edu/akweeds\_ranking\_page.htm">http://akweeds.uaa.alaska.edu/akweeds\_ranking\_page.htm</a>.
- Heffernan, K.E., P.P. Coulling, J.F. Townsend, and C.J. Hutto. 2001. Ranking Invasive Exotic Plant Species in Virginia. Natural Heritage Technical Report 01-13. Virginia Dept. of Conservation and Recreation, Division of Natural Heritage, Richmond, Virginia. 27 pp. plus appendices (total 149 p.).
- Morse, L.E., J.M. Randall, N. Benton, R. Hiebert, and S. Lu. 2004. An Invasive Species Assessment Protocol: Evaluating Non-Native Plants for Their Impact on Biodiversity. Version 1. NatureServe, Arlington, Virginia. http://www.natureserve.org/getData/plantData.jsp
- Randall, J.M., L.E. Morse, N. Benton, R. Hiebert, S. Lu, and T. Killeffer. 2008. The Invasive Species Assessment Protocol: A Tool for Creating Regional and National Lists of Invasive Nonnative Plants that Negatively Impact Biodiversity. Invasive Plant Science and Management 1:36–49
- Warner, Peter J., Carla C. Bossard, Matthew L. Brooks, Joseph M. DiTomaso, John A. Hall, Ann M.Howald, Douglas W. Johnson, John M. Randall, Cynthia L. Roye, Maria M. Ryan, and Alison E. Stanton. 2003. Criteria for Categorizing Invasive Non-Native Plants that Threaten Wildlands. Available online at www.caleppc.org and www.swvma.org. California Exotic Pest Plant Council and Southwest Vegetation Management Association. 24 pp.
- Williams, P. A., and M. Newfield. 2002. A weed risk assessment system for new conservation weeds in New Zealand. Science for Conservation 209. New Zealand Department of Conservation. 1-23 pp.